

**APPLICATION OF GENERALIZED HALLUCINATION THERAPY TO
REDUCE THE LEVEL OF HALUCINOTS IN SCHIZOPHRENIC
PATIENTS AT Dr.RM SOEDJARWADI HOSPITAL, KLATEN**

Fadhilah Intan Pratiwi¹, Sitti Rahma Soleman²

¹Student of the Nursing Profession Program, University of 'Aisyiyah Surakarta

² Lecturers of the Nursing Professional Study Program, 'Aisyiyah University,
Surakarta

Fadhilahip9@gmail.com

ABSTRACT

Background: Hallucinations are false perceptual distortions that occur in maladaptive neurobiological responses, sufferers actually experience sensory distortions as real things and respond to them (Pardede, 2020). Generalist therapy is the application of standard scheduled nursing care that is applied to patients with the aim of reducing the mental nursing problems being handled. (Livana et al., 2020).**Objective:** To find out the results of implementing Generalist Therapy to Reduce the Level of Hallucinations in Schizophrenic Patients at RSJD Dr.RM Soedjarwadi Klaten.**Method:** The method used was a case study research design, the sample used was 2 respondents, the research instrument used was the AHRs observation sheet which was carried out before and after Hallucination Generalist Therapy was carried out.**Results:** Before Generalist Hallucination Therapy Mr. A was in the category of severe hallucinations while Mr. P was in the category of moderate hallucinations. After Generalist Hallucination Therapy Mr. A was in the category of seeding hallucinations while Mr. P was in the mild category.**Conclusion:** There was a decrease in the level of hallucinations after the Hallucination Generalist Therapy was carried out on Mr. A and Mr.P which was carried out for 3 meetings.

Keywords: *Hallucinations, Hallucination Generalist Therapy*