

**THE APPLICATION OF FREE DRAWING THERAPY TO REDUCE THE SIGNS OF
HALUCINOTS IN SCHIZOPHRENIC PATIENTS WITH HALUSCINATIONS
AT RSJD Dr. RM SOEDJARWADI KLATEN
PROVINCE OF CENTRAL JAVA**

Faizah Nur Fajriyati¹, Tri Susilowati²
Fajrifaizah81@gmail.com
University of ' Aisiyiah Surakarta

ABSTRACT

Background rear : Disturbance soul is problem mental health can happen almost on residents all over country in the world . According to *World Health Organization* state schizophrenia is chronic mental disorders And severely affecting 20 million people worldwide _ world (WHO, 2019) Based on existing data at RSJD Dr. RM. Soedjarwadi Province Central Java client data disturbance soul with Schizophrenia on in 2015 there were 751 people , in 2016 there were 853 people , in 2017 there were 981 people , and on year 2018 since month January until February 365 souls . Amount client Schizophrenia at Home Sick Regional Soul Dr. RM. Soedjarwadi Province Central Java on four year final This Keep going experience improvement . Based on background behind problem And the results obtained above , then writer interested For take action nursing about “ Applying Therapy draw free to decline sign symptom hallucinations on on patient schizophrenia with hallucinations at RSJD RM Dr. Soedjarwadi Klaten ” because the most problems happened at Rsjd Dr. Rom. Soedjarwadi Klaten is hallucinations **Purpose :** Knowing results implementation effectiveness therapy activity group : Drawing free to decline sign And symptom hallucinations . **Method :** Study use design *one group pretest posttest*. **Result :** Result application therapy draw free to decline sign And symptom hallucinations on Respondent I experienced decline of the total results 9 to 4 meanwhile respondent II experienced decline from 6 to 2. **Conclusion :** Therapy activity group Draw free can lower sign And symptom hallucinations on patient Hallucinations at RSJD Dr. RM. Soedjarwadi Klaten .

Keywords : _ *Therapy activity group , Draw free , Hallucination*