

**APPLICATION OF DRAWING OCCUPATIONAL THERAPY TOWARDS
THE DECREASE OF AUDITORIAL HALLUCINATIONS IN
SCHIZOPHRENIA PATIENTS AT DR. RM. SOEDJARWADI
RSJD CENTRAL JAVA PROVINCE**

Putri Wulandari¹, Amalia Arifatul Diktina², Wahyu Reknoningsih³

Aisyiyah University of Surakart^{1,2}

RSJD Dr. RM. Soedjarwadi Provinsi Jawa Tengah³

putwulandari166@gmail.com

ABSTRACT

Background: *The most common mental disorder problem is auditory hallucinations. Hallucination sufferers experience inaccuracy in perceiving stimuli or objects that are not actually real, but are considered real.* **Objective:** *Knowing the results of the implementation of occupational drawing therapy on reducing auditory hallucinations in schizophrenia patients at Dr. RM. Soedjarwadi Mental Hospital, Central Java Province.* **Methods:** *Using a case study design, subjects in 2 respondents who were treated at Dr. RM. Soedjarwadi Mental Hospital, Central Java Province were given implementation for 2 consecutive days as much as 1x a day for a duration of 20-30 minutes.* **Results:** *The results were obtained between before and after the implementation of occupational drawing therapy, the condition of both respondents showed a decrease in auditory hallucinations, namely in respondent I from 31 to 17 and respondent II from 25 to 9.* **Conclusion:** *There was a decrease in auditory hallucinations after being given occupational drawing therapy in both respondents.*

Keywords: *Schizophrenia, Hallucinations, Occupational Drawing Therapy*