

## ABSTRAK

Irna Yuliati B2015053 Program Studi DIII Keperawatan STIKES 'Aisyiyah Surakarta	Dosen Pembimbing : 1. Mursudarinah M.Kes 2. Anjar Nurrohmah,.M. Kep
<b>PENERAPAN PIJAT OKSITOSIN UNTUK MEMPERCEPAT INVOLUSI UTERI PADA IBU POST PARTUM DI BPM FINULIA SURAKARTA 2018</b>	
<b>ABSTRAK</b>	
<p><b>Latar Belakang</b> ;Involusi uterus merupakan proses kembalinya uterus ke bentuk semula sebelum hamil dengan berat 60 gram. Prevelensi perdarahan data di klinik BPM Finulia Surakarta pada bulan januari sampai desember tahun 2017 di dapatkan angka persalinaan pada tahun sebanyak 30 orang 5 diantaranya mengalami perdarahan akibat melemahnya kontraksi uterus. Involusi uteri pada pasien bisa diatasi dengan terapi farmakologis dan nonfarmakologis, salah satu terapi nonfarmakologis adalah pijat oksitosin. Pijat oksitosin adalah Pijat oksitosin adalah tindakan yang dilakukan pada ibu berupa pijatan pada punggung ibu untuk meningkatkan pengeluaran hormon oksitosin, pijat oksitosin dilakukan 2 kali sehari selama 2-3 menit 7 hari berturut-turut</p> <p><b>Tujuan</b>;Mendiskripsikan hasil penerapan teknik pijat oksitosin untuk mempercepat involusi uteri pada ibu postpartum</p> <p><b>Metode Penelitian</b> ;Menggunakan metode diskriptif dengan dengan mengobservasi pada subyek penelitian. Pengukuran tinggi fundus uteri diukur dengan metline lalu dicatat sebelum dan sesudah penerapan pijat oksitosin.</p> <p><b>Hasil</b>;Pengukuran tinggi fundus uteri sebelum dilakukan pijat oksitosin 13 cm dan 14 cm, sedangkan tinggi fundus uteri sesudah dilakukan pijat oksitosin tidak teraba dan 3 cm</p> <p><b>Kesimpulan</b>;Terdapat penurunan tinggi fundus uteri setelah dilakukan pijat oksitosin.</p>	
<b>Kata Kunci</b> : <i>Postpartum, Involusi Uteri, Pijat Oksitosin</i>	

## ABSTRACT

Irna Yuliati B2015053 Program Studi DIII Keperawatan STIKES 'Aisyiyah Surakarta	Counsellor : 1. Mursudarinah M.Kes 2. Anjar Nurrohmah,.M. Kep
<b><i>The Influence Of Oxytocin Massage With Uterus Involution Mother's Post Partum In BPM Finulia Surakarta 2018</i></b>	
<b>ABSTRACT</b>  <i><b>Background</b> ; Uterine involution is a process of return of the uterus to form before pregnancy weighing 60 grams. The prevalence of data bleeding at BPM clinic Finulia Surakarta from January to December 2017 was found in the number of pers in the year as many as 30 people 5 of whom suffered bleeding due to weakening uterine contractions. The uterine involution of the patient can be treated with pharmacological and nonpharmacological therapy, one of the nonpharmacological therapies being the oxytocin massage. Massage oxytocin is a massage of oxytocin is an action performed on the mother in the form of massage on the mother's back to increase the expenditure of the hormone oxytocin, oxytocin massage done 2 times a day for 2-3 minutes 7 days in a row.<b>Objective</b>; Describes the application of oxytocin massage techniques to accelerate uterine involution in postpartum mother. <b>Research Methods</b>; Using the descriptive method by observing on the subject of the study. Measurement of uterine fundus height was measured by metline and then recorded before and after the application of oxytocin massage. <b>Results</b>; Measuring the height of the uterine fundus prior to oxytocin massage 13 cm and 14 cm, while the height of the uterine fundus after the invisible oxytocin massage and 3 cm. <b>Conclusion</b>; There is a decrease in uterine fundus height after oxytocin massage.</i>	
<b>Keywords:</b> <i>Post Partum, Uterus Involution, Oxytocin Massage</i>	