

**THE IMPLEMENTATION TOPICAL WITH BREAST MILK  
TOWARD THE LENGTH TIME OF UMBILICAL CORD RELEASE  
ON NEWBORN BABY AT THE MOJOLABAN HEALTH CENTER**

Elysa Dwi Sholiqah, Siti Fatmawati, Irma Mustika Sari

[Elisholeha1@gmail.com](mailto:Elisholeha1@gmail.com)

Diploma III Nursing Study Program  
STIKES Aisyiyah of Surakarta

**ABSTRACT**

**Background:** Indonesia is the one of countries that has a high infant mortality rate. Neonatal mortality is caused due to the umbilical cord infections. Infection of the umbilical cord care can cause tetanus neonatrum in infants, the main cause of tetanus neonatrum because of poor cord care. Whereas, the umbilical cord care becomes the thing that requires special attention. That is an effort to prevent the umbilical core infection, which is need cleanliness and hygienically. And one of the umbilical cord care uses topical with breast milk. **Objective:** this study aimed to describe umbilical cord care with breast milk for the length of time of umbilical cord release. **Method:** the research was conducted by using descriptive qualitative method on case study. The sample of the research are two respondents. While the instrument of the research used observation sheets. **Results:** the umbilical cord release using topical with breast milk loose on day 4 of the fast category and day 6 of the normal category. The average time of umbilical cord release using topical with breast milk was 4-6 days.

**Keywords:** *The umbilical cord, topical with breast milk*