

***APPLICATION OF MUROTTAL THERAPY TO CHANGE
SCHIZOPHRENIC CLIENT VIOLENT BEHAVIOR
AT RSJD Dr. RM. SOEDJARWADI KLATEN
PROVINCE OF CENTRAL JAVA***

Siti Putri Yuliana¹, Sitti Rahma Soleman², Wahyu Reknoningsih³
Pyuliana45@gmail.com
Universitas 'Aisyiyah Surakarta
ABSTRACT

Background; The risk of violent behavior is a response to the stressor faced by someone who is shown by violent behavior both to himself and to other people and the environment both verbally and non-verbally. One way to reduce the impact of violent behavior is by giving murottal therapy. **Objective;** To find out the results of implementing the Murottal Therapy Application for Changes in the Violent Behavior of Schizophrenic Clients at RSJD Dr. RM. Soedjarwadi, Central Java Province. **Method;** This type of research is a case study using descriptive research methods. **Results;** The score for violent behavior change for 2 respondents before the murottal therapy was carried out was at the intensive stage score 2, which means that at the moderate behavior change stage where client 1 got a score of 11 while client 2 got a score of 14. The score for the change in violent behavior for 2 respondents after murottal therapy was at the stage score intensive 3 which means at the stage of mild behavior change where client 1 gets a score of 8 while client 2 gets a score of 9. There are differences in changes in violent behavior in respondents with a risk of violent behavior before and after the murottal therapy intervention. **Conclusion;** There are differences in changes in violent behavior before and after murottal therapy in patients at risk for violent behavior.

Keywords: Decrease, Risk of Violent Behavior, Murottal Therapy